

LETTER • **OPEN ACCESS**

Black truffle winter production depends on Mediterranean summer precipitation

To cite this article: Ulf Büntgen *et al* 2019 *Environ. Res. Lett.* **14** 074004

View the [article online](#) for updates and enhancements.

Recent citations

- [The crop of desert truffle depends on agroclimatic parameters during two key annual periods](#)
Alberto Andriano *et al*

Environmental Research Letters



LETTER

OPEN ACCESS

RECEIVED

6 December 2018

REVISED

8 April 2019

ACCEPTED FOR PUBLICATION

11 April 2019

PUBLISHED

1 July 2019

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Black truffle winter production depends on Mediterranean summer precipitation

Ulf Büntgen^{1,2,3,4} , Daniel Oliach^{5,6,7}, Fernando Martínez-Peña^{8,9}, Joaquin Latorre⁸, Simon Egli² and Paul J. Krusic^{1,10}¹ Department of Geography, University of Cambridge, Downing Place, CB2 3EN, United Kingdom² Swiss Federal Research Institute WSL, Zürcherstr 111, 8903 Birmensdorf, Switzerland³ Global Change Research Centre (CzechGlobe), Bělidla 986/4a, Brno 603 00 Brno, Czech Republic⁴ Department of Geography, Faculty of Science, Masaryk University, Kotlářská 2, 613 00 Brno, Czech Republic⁵ Centre Tecnològic Forestal de Catalunya, Crta. Sant Llorenç de Morunys km 2, E-25280 Solsona, Spain⁶ Forest Bioengineering Solutions S. A., Crta. Sant Llorenç de Morunys km 2, E-25280 Solsona, Spain⁷ Universitat de Lleida-AGROTECNIO. Dept. de Producció Vegetal i Ciència Forestal. Av. Rovira Roure, 191, E-25198 Lleida, Spain⁸ European Mycological Institute EGTC-EMI, E-42003 Soria, Spain⁹ Agrifood Research and Technology Centre of Aragon CITA, Montañana 930, E-50059 Zaragoza, Spain¹⁰ Department of Physical Geography, Stockholm University, SE-106 91 Stockholm, SwedenE-mail: ulf.buentgen@geog.cam.ac.uk**Keywords:** climate change, fungal ecology, economic sustainability, irrigation practices, Mediterranean drought, truffle production, *Tuber melanosporum*Supplementary material for this article is available [online](#)

Abstract

The unprecedented price inflation of Black truffles, recently exceeding 5000 Euro kg⁻¹ (in Zurich), is a combined result of increasing global demands and decreasing Mediterranean harvests. Since the effects of long-term irrigation and climate variation on symbiotic fungus-host interaction and the development of belowground microbes are poorly understood, the establishment and maintenance of truffle plantations remains a risky venture. Using 49 years of continuous harvest and climate data from Spain, France and Italy, we demonstrate how truffle production rates, between November and March, significantly rely on previous June–August precipitation totals, whereas too much autumnal rainfall affects the subsequent winter harvest negatively. Despite a complex climate-host-fungus relationship, our findings show that southern European truffle yields can be predicted at highest probability ($r = 0.78$, t -stat = 5.645, prob = 0.000 01). Moreover, we demonstrate the reliability of national truffle inventories since 1970, and question the timing and dose of many of the currently operating irrigation systems. Finally, our results suggest that Black truffle mycorrhizal colonization of host fine roots, the sexualisation of mycelium, and the formation of peridium are strongly controlled by natural summer rainfall. Recognising the drought-vulnerability of southern Europe's rapidly growing truffle sector, we encourage a stronger liaison between farmers, politicians and scientists to maintain ecological and economic sustainability under predicted climate change in the Mediterranean basin.

Introduction

Approximately 16 km² of arable land in northeastern Spain and southern France are transformed each year into new plantations of the (Périgord) Black truffle (*Tuber melanosporum* Vittad, an Ascomycota; herein-after 'truffle'), with slightly smaller units in north-central Italy. This booming industry substantially contributes to rural economies and cultural identity

(Samils *et al* 2008, Büntgen *et al* 2017). An estimated 110 000 kg yr⁻¹ of truffles that are growing in wild habitats and ~40 000 ha of scattered plantations in Spain, France and Italy generate ~50 million Euro annually (Oliach *et al* 2019). The rapidly growing and wide-ranging economic sector includes the production of mycorrhized plants in nurseries, the harvest of wild and cultivated truffles, truffle dog training, the marketing of fresh and processed truffles, the

transformation of truffles into secondary products, mycotourism (i.e. truffle-tourism), mycological gastronomy, interdisciplinary research, and producer extension services (Büntgen *et al* 2017). The total asset, in France alone, has been estimated at ~ 67 million Euro yr^{-1} . With an increasing trend, cultivated fruitbodies already account for up to 80% of all commercially traded truffles (Murat 2015, Reyna and Garcia-Barreda 2014). Due to the hidden belowground life-cycle of this iconic culinary species (Trappe and Claridge 2010), and its complex host interaction (Büntgen *et al* 2015), as well as potential direct and indirect climatic (e.g. precipitation and temperature, respectively) impacts (Büntgen *et al* 2012, Molinier *et al* 2013, Le Tacon *et al* 2014, 2016, Thomas and Büntgen 2019, Baragatti *et al* 2019), truffle cultivation in southern Europe is still associated with high ecological and economic risks. Although many plantations are now irrigated, an increase in the frequency and intensity of Mediterranean summer droughts is expected to affect both, the quality and quantity of the subsequent truffle winter harvest (Büntgen *et al* 2012, Thomas and Büntgen 2019). This is particularly alarming since warming in southern Europe is predicted to exceed global rates by 25% (Cramer *et al* 2018), notably with summer temperatures rising at a pace 40% larger than the worldwide mean (Lionello and Scarascia 2018). This trend will be associated with more heatwaves and a reduction in summer precipitation of around 10%–15% over the Mediterranean truffle producing regions (Fischer and Schär 2010, Büntgen *et al* 2012, Jacob *et al* 2014, Vautard *et al* 2014, Thomas and Büntgen 2019).

Despite a putative sensitivity bias, due to a substantial increase in irrigation intensity and refined cultivation practices during the past years (Olivera *et al* 2014a, 2014b, Oliach *et al* 2019), as well as the common belief that national truffle production data are very noisy because of uncoordinated trading and often-unofficial marketing (Reyna 2012), in addition to many other biases (Le Tacon *et al* 2014, Baragatti *et al* 2019), this study aims to understand the dependency of truffle productivity on rainfall variability. To reconstruct the effects of Mediterranean climate change on truffle harvest, we collected annual estimates of the fungus' fruitbody yield from Spain, France and Italy between 1970 and 2017/18, and compared these values with monthly resolved and spatially explicit temperature and precipitation indices. Time series analyses, spatial field correlation coefficients and a suite of calibration-verification trials were applied to quantify the relationship between truffle production and climate variation at different spatiotemporal scales.

Data and methods

Three continuous, 49 year-long records of the approximate annual truffle production from southern Europe's main truffle producing regions in northeastern Spain, southern France and northcentral Italy is analysed and compared against high-resolution, gridded climate indices of the same three regions. Though still associated with wide uncertainties (see discussion below), data from the national harvest inventories represent a substantial update in the number of years studied from initial 37 (1970–2005/6) in Büntgen *et al* (2012) to the current 49 years from 1970 to the latest complete truffle harvest between November 2017 and March 2018 (supplementary online material is available online at stacks.iop.org/ERL/14/074004/mmedia). Information on the winter truffles harvest from northeastern Spain and southern France was compiled by the national Truffle Grower Associations and the Groupement European Tuber (Courvoisier 1992, Callot 1999, Reyna 2012, Oliach *et al* 2019). The French Ministry of Agriculture gathered yields from across France until 1988, whereas data afterwards are restricted to the most important markets, from which the French National Truffle Grower Association calculated the nationwide harvest. Truffle information from northcentral Italy was collected and published by the National Institute for Statistics until 1990, and afterwards from the Groupement European Tuber (Oliach *et al* 2019). While allowing year-to-year variability to be analysed, the limited resolution of the truffle data does not reflect any intra-seasonal changes.

For comparison against regional and Mediterranean-wide climate variability, monthly and spatially resolved gridded temperature means and precipitation totals were extracted from the E-OBS v8.0 reanalyses dataset (Haylock *et al* 2008). In addition to European-scale field correlation analyses (as commonly applied in high-resolution (paleo)climatic studies through the KNMI server; <http://climexp.knmi.nl/>), regional climate indices were derived by averaging the E-OBS grid cells over 41–42 °N and 2–0 °W for northeastern Spain, over 44–45 °N and 3–5 °E for southern France, and over 44–45 °N and 10–12 °E for northcentral Italy. The normalized grid cell averages (i.e. Z-scores with a mean of zero and a standard deviation of one over 1970–2018), climatically representative of the truffle producing regions in each country, were used as predictors of similarly normalized truffle production (mean of 0 and STDV of 1). Pearson's correlation coefficients were used to determine those months (i.e. each individual monthly value of the year of truffle growth) and/or seasons (i.e. averages of two or more consecutive months prior to truffle harvest) when regional climate indices are significantly correlated ($p < 0.05$) with truffle harvests (see supplementary online material for details). To test the statistical robustness and temporal stability of the relationship between monthly

climate and fruitbody production this experiment is performed three times for each region, once for the first half of the truffle and climate data's common period (1970–1993), once for the second half (1994–2017), and once for the full period (1970–2017). Dividing the period shared by both the harvest and precipitation data into an early-period and a late-period, and using the linear model estimates derived from the information in one to verify the unused values in the second, and visa-versa, is called split period calibration/verification. This method of establishing a robust linear model between an instrumental quantity and an independent time series of measurements is common practice in dendroclimatology (Esper *et al* 2016).

The skill with which modelled estimates of truffle winter production, derived from climate values in the calibration period, replicates the observed variance of production in the validation period, is expressed by the performance of the coefficient of efficiency (CE) and reduction of error (RE) statistics. Both RE and CE are measures of the shared variance between actual and estimated values (with CE being the more rigorous statistic). Positive values of RE and CE suggest the model has predictive skill (Frits 1976, Cook *et al* 1994). As often used in high-resolution palaeoclimatology, the Durbin–Watson (DW) statistic assesses temporal stability in the calibration models (DW; Durbin and Watson 1951). DW tests for lag-1 autocorrelation in model residuals. A DW value > 1.00 , for $n = 47$, represents an acceptable degree of first-order autocorrelation in the residuals ($p < 0.05$).



To guard against inflated correlations due to covariance in trend (low-frequency), all further modelling experiments are performed on both first-differences (FD; high-frequency), and undifferenced (UD; actual values) data. The calibration and verification exercises, using monthly and seasonal precipitation totals as predictors of truffle production, are the final step before accepting any model's hind or forecast ability. These experiments are performed on the FD transformations of seasonal precipitation and winter production of the truffle, as well as their original UD values. Once again, the classical split period approach is applied independently for each of the truffle producing regions. When both periods produce verifiable estimates, significantly correlated with only positive error reduction, the relationship between predictor and predictand is considered robust, and a reconstruction or prediction may be performed.

Results

None of the monthly and seasonal temperature means reveal significantly ($p < 0.05$) positive relationships with any of the regional or Mediterranean-wide truffle winter yields (figure 1). However, Pearson's correlation coefficients between truffle winter production

and monthly summer precipitation reveal significantly ($p < 0.05$) positive values at both the regional and sub-Mediterranean scales. Despite current irrigation efforts, truffle harvests in northeastern Spain, southern France and northcentral Italy exhibit their highest correlations with rainfall in slightly different summer months (figure 1). Moreover, precipitation totals between October and November have significant ($p < 0.05$) negative effects on the subsequent fungal yield. The total truffle harvest of all three regions is significantly positively correlated with total June–August precipitation (figures 1, 2; table 1). This association is confirmed by DW statistics of 1.1056, 1.1569 and 1.3795 for Spain, France and Italy, respectively. Over the full 1970–2017/18 period (figure 2(A)), the Spanish harvest has the highest correlation with summer precipitation ($r = 0.68$), followed by France and Italy ($r = 0.59$). All three regions display a long-term decline in both, summer rainfall and winter truffle yield, from the mid-1970s until around 2000. While there is a sharp rise in Spanish summer precipitation and truffle winter production from 2012 to present, the French data present a much slower, though continuous increase since around 2003. This positive trend is less distinct in Italy, where truffle production peaked between 2012 and 2014. Most surprising is the strong dependency of truffle winter production on previous summer precipitation in Spain since 1994 ($r = 0.75$), the driest region and period in which irrigation is most intense. Another surprising result is the high agreement between truffle data from Spain and those from France and Italy ($r = 0.52$ and 0.47), which is not mirrored by their corresponding precipitation records ($r = 0.48$ and 0.17). In addition to the temporally stable association between truffle winter production and June–August precipitation (figure 2(A)), European-wide field correlations of each of the three production regions exhibit remarkably strong spatial coverage of explained summer rainfall variability (figure 2(B)). The highest correlations are again found over Spain, from the Iberian System in the south to the foothills of the Pyrenees in the north, followed by two clusters in southern France, west and east of the Rhone Valley, and along a north-to-south transect in Italy between the Po Valley in the north and the Apennine Mountains in the south. Another interesting finding are the negative correlations between truffle production and precipitation between September/October and November (figure 1), which are distinct in all three countries.

Full period modelling using the optimal seasonal precipitation averages as a predictor of truffle winter harvest reveals statistically significant solutions (figure S1), the most robust of which is that for the Spanish harvest (table 1). Time series analysis of model residuals (figure S1), in all instances, shows there is still unaccounted persistence manifest by the rather low DW statistics. This suggests that, in addition to the



variation accounted for by precipitation there is in fact an additional, unaccounted, factor affecting fruitbody production that is transient in nature (see Discussion below).

The best prediction model is that which explains the Spanish truffle winter production (figure S2). The French model is also arguably acceptable, however, it is clear that the Italian model, though a robust

Table 1. Linear regression models for regional precipitation totals and truffle yields. Spanish, French and Italian Summer (June–August) precipitation totals as a predictor of truffle production in the corresponding countries between 1970 and 2017 explains 46.22%, 28.43% and 25.62% variance, respectively.

	Corr.	t-stat	Prob.	RSQ	Cum. RSQ	Adj. RSQ	Adj. RE	AIC
Spain	0.680	6.288	0.0000	0.462	0.462	0.451	0.439	−25.51
France	0.533	4.274	0.0001	0.284	0.284	0.269	0.254	−11.79
Italy	0.506	3.981	0.0003	0.256	0.256	0.240	0.225	−9.94

Table 2. Calibration/verification models for Mediterranean precipitation totals and truffle yields. Combined Spanish, French and Italian Summer (June–August) precipitation totals as a predictor of total Mediterranean truffle production for two early/late split periods. All statistics are calculated for normalized undifferenced and pre-whitened values (first-differenced), with positive reduction of error (RE) and coefficient of variation (CE) values suggesting strong verification results (see tables S1–S3 for the regional calibration/verification models). The verification period, RE is an implementation of Allen’s PRESS statistic (Allen 1974), which employs a leave-one-out calculation between actual and estimated values, similar to the leave-one-out iterative calculations commonly used in cross-validation tests.

Calibration period results (undifferenced data 1970–1993)									
Pearson	Prob.	Robust	Prob.	Spearman	Prob.	RE	CE	MedRE	MedCE
0.606	0.001	0.601	0.001	0.606	0.000	0.368	0.368	0.365	0.337
Calibration period results (1st-differenced data 1970–1993)									
Pearson	Prob.	Robust	Prob.	Spearman	Prob.	RE	CE	MedRE	MedCE
0.607	0.001	0.432	0.001	0.451	0.001	0.324	0.324	0.269	0.242
Verification period results (undifferenced data 1994–2017)									
Pearson	Prob.	Robust	Prob.	Spearman	Prob.	RE	CE	MedRE	MedCE
0.551	0.000	0.525	0.000	0.416	0.000	0.475	0.149	0.461	0.091
Verification period results (1st-differenced data 1994–2017)									
Pearson	Prob.	Robust	Prob.	Spearman	Prob.	RE	CE	MedRE	MedCE
0.776	0.000	0.768	0.000	0.701	0.000	0.517	0.506	0.448	0.438


predictor of annual variation, lacks the most credibility in predicting low-frequency behaviour (tables S1–S3). Repeating the calibration and verification experiment using the averages of all three regional summer precipitation totals, and the average of the three national truffle winter production records, produces another verifiably robust model (figure S3). All verification statistics of RE and CE are positive for each period, and at both the high and low-frequency domains (table 2). This strong sub-Mediterranean dependency of truffle winter production to previous summer precipitation is mirrored in the spatial correlation fields of the averaged data (figure 3). When calculated over the full period 1970–2017/18, significantly ($p < 0.05$) positive correlations cover most of the truffle producing regions in all three countries (figure 3(A)), with an overall increase in this relationship towards present (figures 3(B), (C)).

Discussion

To some degree, our results call into question the timing and dose of those irrigation systems that already operate. This is particularly the case for north-eastern Spain since the 1990s, which represents the driest period and region of this study. Although the

current irrigation prescription seems ineffective, we argue that a simple increase of the amount of water might not be helpful (Bonet *et al* 2006, Olivera *et al* 2014a, Büntgen *et al* 2015), since the formation and maturation of truffle fruitbodies is likely enhanced by periodic drought-stress (Garcia-Barreda *et al* 2019). Since temporally adjustable belowground watering systems might be more efficient than traditional aboveground sprinklers, such techniques could reduce the burden of current water-use allowances, which are predicted to become more restrictive as the frequency and severity of Mediterranean summer droughts increases (Fischer and Schär 2010, Trnka *et al* 2018). While deficit irrigation might be an alternative (Ferreira and Soriano 2006, Sears *et al* 2018), it requires understanding of the fungus’ full lifecycle (Baragatti *et al* 2019).


Consistent with previous findings (Gallot 1999, Büntgen *et al* 2012, 2015, Le Tacon *et al* 2014, 2016, Thomas and Büntgen 2019, Baragatti *et al* 2019), our results highlight the importance of summer precipitation for truffle winter production. High precipitation totals and low temperature means between June and August are expected to stimulate (a) mycorrhizal colonization of host fine roots, (b) formation and sexualisation of mycelium, and (c) development of peridium. We further assume that the truffles’



associated tree partners not only provide an important carbon pool, but also act as the principal source of water in dry periods during fruitbody formation and maturation. We speculate that the host plants, water stressed themselves, may possibly also provide compensation to the fungi during drought spells through increased hydraulic lifting. A better understanding of the potential hydraulic redistribution of soil water by direct nocturnal water transfer from host trees to their mycorrhizal symbionts is, however, needed (Querejeta *et al* 2003, Warren *et al* 2008). Favouring environments of contrasting drought-stress (Garcia-Barreda *et al* 2019), truffles can survive dry periods of up to 30 d (Ricard *et al* 2003). Our own observations of *T. aestivum* in Switzerland and southern Germany suggest that fruitbodies can mature in totally dried-out soils, where other epigeous ectomycorrhizal species have already stopped fruiting. Another interesting finding is the inverse relationship between winter truffle yields and precipitation totals in October and November (figure 1), which indicates that wetter and cooler

autumns negatively affect the subsequent truffle harvest. Furthermore, we did not find any significant positive effects of monthly and/or seasonal temperature means (figure 1), which supports the idea that truffles can grow under much cooler (or warmer) conditions than previously thought (Thomas and Buntgen 2017). It is important to note that June–August precipitation totals and temperature means in Spain, France and Italy are significantly negatively ($p < 0.05$) correlated ($r = -0.46$, -0.47 and -0.62 , respectively). While the obtained truffle-climate relationships appear most reasonable in a myco-bio/ecological perspective (Buntgen *et al* 2015), the role of host plants is largely unknown (Buntgen and Egli 2014), and some bias may emerge from imprecise meteorological measurements that were aggregated over broad spatiotemporal scales rather than reflecting the environmental conditions of the exact locations and periods of truffle growth.

In addition to the direct negative effects of a dryer future on the growth and development of truffles and



their hosts—similar to other ectomycorrhizal fungi (Köhler *et al* 2018), there are several indirect, temperature-induced, factors (Baragatti *et al* 2019), such as wildfires, pathogens and diseases (Thomas and Büntgen 2019), as well as phenological mismatch in trophic interactions that may disrupt current ecological systems (Renner and Zohner 2018), and cause economic damage. A longer fire season combined with more frequent large fires is expected as a result of increasing summer temperatures, drought and land-use changes (Khabarov *et al* 2016, Ruffault *et al* 2016). Forest fires not only kill trees but also impact soil chemistry, which affects ectomycorrhizal fungal communities (Mediaviella *et al* 2017). Similarly, the wide range of insect pests and destructive pathogens, such as *Phytophthora cinnamomi* that feed on oaks, are expected to expand their distribution under warmer winter temperatures (Bergot *et al* 2004, Barredo *et al* 2015). In addition, insect pest may also directly affect truffle fruitbodies and thus pose a serious threat to the emerging industry (Rosa-Gruszecka *et al* 2017), because warming increases both population growth and metabolic rates of insects (Deutsch *et al* 2018), and even small larvae infestations already cause large damage. Almost ironically, a warmer and dryer future implies more flood hazards (Cramer *et al* 2018), which can trigger massive surface erosion and sediment relocation, associated with reductions in the richness and abundance of ectomycorrhizal fungi (Barnes *et al* 2018).

Conclusions

This study shows that inventories of truffle yield from Spain, France and Italy, rather than reflecting mainly noise, are reliable since 1970, and that winter truffle harvests significantly depend on previous summer rainfall, whereas too much autumnal precipitation has negative effects. Our findings question the timing and dose of the existing irrigation systems, and call for both management and conservation action to mitigate a multitude of unprecedented ecological and economic risks under predicted climate change. The various threats might be particularly severe for rural Mediterranean cultivators that are most vulnerable to a warmer and dryer future (Büntgen *et al* 2017, Cramer *et al* 2018).

Ultimately, we provide a robust tool for predicting sub-Mediterranean truffle winter production from previous summer precipitation at highest probability. The degree of statistical significance afforded by our model ($r = 0.78$, $t\text{-stat} = 5.645$, $\text{prob} = 0.000\,01$) rivals that of the best high-resolution climate proxy records (Esper *et al* 2016), for instance. Considering, the number of environmental factors not accounted for in this linear model, such as the inverse relationship with summer temperature (Thomas and Büntgen 2019, Baragatti *et al* 2019), it is remarkable to find summer precipitation alone can explain 36.76% of the subsequent truffle winter production (table 2; figure 4). If handled responsibly, this information can help stabilize production and pricing from regional to international scales, thereby contributing to the maintenance of sustainable harvests and markets. The likelihood to forecast truffle production from summer to

winter, however, does not enable long-term projections since host density and irrigation intensity in plantations can (and should) be adapted to changing environmental conditions. Since a drought-induced collapse of the system would also trigger biodiversity losses, a critical review of the current plantation practices seems timely and calls for a vibrant liaison between academia, policy and economy at local to international levels. Finally, we hope that our study will stimulate more detailed work to explore the species' full lifecycle, such as yearlong, fine-scale excavating technique from archaeology as a new approach in ectomycorrhizal research to gain unique insights into the hidden belowground truffle kingdom.

Acknowledgments

This work was supported by the European Mycological Institute (EMI), as well as the WSL-internal project 'Black Dynamite'. UB received funding from the project 'SustES - Adaptation strategies for sustainable ecosystem services and food security under adverse environmental conditions' (CZ.02.1.01/0.0 /0.0/16_019/0000797). DO received support from the Secretaria d'Universitats i Recerca del Departament d'Economia i Coneixement de la Generalitat de Catalunya through the program of Doctorats Industrials. The authors are very grateful to The European Truffle Group (GETT), entity composed by French, Spanish and Italian growers' federations, for its considerable help with truffle production data.

ORCID iDs

Ulf Büntgen  <https://orcid.org/0000-0002-3821-0818>

References

- Allen D M 1974 The relationship between variable selection and data augmentation and a method for prediction *Technometrics* **16** 125–7
- Baragatti M, Grollemund P-M, Montpied P, Dupouey J-L, Gravier J, Murat C and Le Tacon F 2019 Influence of annual climatic variations, climate changes, and sociological factors on the production of the périgord black truffle (*Tuber melanosporum* Vittad.) from 1903–1904 to 1988–1989 in the Vaucluse (France) *Mycorrhiza* **29** 113–25
- Barnes C J, van der Gast C J, McNamara N P, Rowe R and Bending G D 2018 Extreme rainfall affects assembly of the root-associated fungal community *New Phytol.* **220** 1172–84
- Barredo J I, Strona G, Rigo D, Caudullo G, Stancanelli G and San-Miguel-Ayán J 2015 Assessing the potential distribution of insect pests: case studies on large pine weevil (*Hylobius abietis* L) and horse-chestnut leaf miner (*cameraria ohridella*) under present and future climate conditions in European forests *EPPO Bull.* **45** 273–81
- Bergot M, Cloppet E, Pérarnaud V, Déqué M, Marçais B and Desprez-Loustau M-L 2004 Simulation of potential range expansion of oak disease caused by *phytophthora cinnamomi* under climate change *Glob. Change Biol.* **10** 1539–52
- Bonet J A, Fischer C R and Colinas C 2006 Cultivation of black truffle to promote reforestation and land-use stability *Agron. Sustain. Dev.* **26** 69–76
- Büntgen U and Egli S 2014 Breaking new ground at the interface of dendroecology and mycology *Trends Plant Sci.* **19** 613–4
- Büntgen U, Egli S, Camarero J J, Fischer E M, Stobbe U, Kauserud H, Tegel W, Sproll L and Stenseth N C 2012 Drought-induced decline in Mediterranean truffle harvest *Nat. Clim. Change* **2** 827–9
- Büntgen U, Latorre J, Egli S and Martínez-Peña F 2017 Socio-economic, scientific and political benefits of mycotourism *Ecosphere* **8** e01870
- Büntgen U *et al* 2015 Long-term irrigation effects on Spanish holm oak growth and its black truffle symbiont *Agric. Ecosyst. Environ.* **202** 148–59
- Callot G 1999 *La truffe, la terre, la vie* (Paris: Ed. INRA)
- Cook E R, Briffa K R and Jones P D 1994 Spatial regression methods in dendroclimatology: a review and comparison of two techniques *Int. J. Climatol.* **14** 379e402
- Courvoisier M 1992 *Bilan de la filière truffe en France* (Paris: Ctifl) 119
- Cramer W *et al* 2018 Climate change and interconnected risks to sustainable development in the Mediterranean *Nat. Clim. Change* **8** 972–80
- Deutsch C A, Tewksbury J J, Tigchelaar M, Battisti D S, Merrill S C, Huey R B and Naylor R L 2018 Increase in crop losses to insect pests in a warming climate *Science* **361** 916–9
- Durbin J and Watson G S 1951 Testing for serial correlation in least squares regression *Biometrika* **38** 159–78
- Esper J *et al* 2016 Ranking of tree-ring based temperature reconstructions of the past millennium *Quat. Sci. Rev.* **145** 134–51
- Fereres E and Soriano M A 2006 Deficit irrigation for reducing agricultural water use *J. Exp. Bot.* **58** 147–59
- Fischer E M and Schär C 2010 Consistent geographical patterns of changes in high-impact European heatwaves *Nat. Geosci.* **3** 398–403
- Fritts H C 1976 *Tree-Rings and Climate* (London, New York, San Francisco: Academic) p 576
- García-Barreda S, Sánchez S, Marco P and Serrano-Notivol R 2019 Agro-climatic zoning of Spanish forests naturally producing black truffle *Agric. Forest Meteorol.* **296** 231–8
- Haylock M R, Hofstra N, Klein Tank A M G, Klok E J, Jones P D and New M 2008 A European daily high-resolution gridded data set of surface temperature and precipitation for 1950–2006 *J. Geophys. Res.* **113** D20119
- Jacob D *et al* 2014 EURO-CORDEX: new high-resolution climate change projections for European impact research *Reg. Environ. Change* **14** 563–78
- Khabarov N, Krasovskii A, Obersteiner M, Swart R, Dosio A, San-Miguel-Ayán J, Durrant T, Camia A and Migliavacca M 2016 Forest fires and adaptation options in Europe *Reg. Environ. Change* **16** 21–30
- Köhler J, Yang N, Pena R, Raghavan V, Polle A and Meier I C 2018 Ectomycorrhizal fungal diversity increases phosphorus uptake efficiency of European beech *New Phytol.* **220** 1200–10
- Le Tacon F, Marçais B, Courvoisier M, Murat C and Becker M 2014 Climatic variations explain annual fluctuations in French Périgord black truffle wholesale markets but does not explain the decrease in black truffle production over the last 48 years *Mycorrhiza* **24** 115–25
- Le Tacon F *et al* 2016 Certainties and uncertainties about the life cycle of the périgord black truffle (*Tuber melanosporum* Vittad.) *Ann. Forest Sci.* **73** 105–17
- Lionello P and Scarascia L 2018 The relation between climate change in the Mediterranean region and global warming *Reg. Environ. Change* **18** 1481–93
- Mediavilla O, Hernández-Rodríguez M, Olaizola J, Santos-del-Blanco L, Oria-de-Rueda J A and Martín-Pinto P 2017 Insights into the dynamics of boletus edulis mycelium and fruiting after fire prevention management *Forest Ecol. Manage.* **404** 108–14

- Molinier V, Bouffaud M L, Caste T, Mounier A, Colombet A, Recorbet G, Frochot H and Wipf D 2013 Monitoring the fate of a 30-year-old truffle orchard in Burgundy: from *Tuber melanosporum* to *Tuber aestivum* *Agroforestry Syst.* **87** 1439–49
- Murat C 2015 Forty years of inoculating seedlings with truffle fungi: past and future perspectives *Mycorrhizal* **25** 77–81
- Oliach D *et al* 2019 Las trufas y las turmas ed S-G Mariola, C Rafael and B J Antonio *Los Productos Forestales no Madereros en España: Del Monte A La Industria. Monografías INIA: Seria Forestal* (Madrid: Ministerio de Economía Industria y Competitividad)
- Olivera A, Bonet J A, Oliach D and Colinas C 2014a Time and dose of irrigation impact *Tuber melanosporum* ectomycorrhiza proliferation and growth of *quercus ilex* seedling hosts in young black truffle orchards *Mycorrhiza* **24** 73–8
- Olivera A, Bonet J A, Palacio L and Colinas C 2014b Adequate weed suppression strategies can promote *T. melanosporum* mycelial expansion and seedling growth in newly established black truffle plantations *Ann. Forest Sci.* **71** 495–504
- Querejeta J I, Egerton-Warburton L M and Allen M F 2003 Direct nocturnal water transfer from oaks to their mycorrhizal symbionts during severe soil drying *Oecologia* **134** 55–64
- Renner S S and Zohner C M 2018 Climate change and phenological mismatch in trophic interactions among plants, insects, and vertebrates *Annu. Rev. Ecol. Evol. Syst.* **49** 165–82
- Reyna S 2012 *Truficultura Fundamentos y Técnicas* 2nd edn (Madrid: Mundi-Prensa)
- Reyna S and Garcia-Barreda S 2014 Black Truffle cultivation: a global reality *Forest Syst.* **23** 317–28
- Ricard J M, Bergounoux F, Callot G, Chevalier G, Olivier J M, Pargney J C and Sourzat P 2003 P. *La Truffe, Guide Technique de Trufficulture* (Paris: CTIFL)
- Rosa-Gruszecka A *et al* 2017 Insect-truffle interactions—potential threats to emerging industries? *Fungal Ecol.* **25** 59–63
- Ruffault J, Moron V, Trigo R M and Curt T 2016 Objective identification of multiple large fire climatologies: an application to a Mediterranean ecosystem *Environ. Res. Lett.* **11** 075006
- Samils N, Olivera A, Danell E, Alexander I, Fischer C R and Colinas C 2008 The socioeconomic impact of truffle cultivation in rural Spain *Econ. Bot.* **62** 331–40
- Sears L, Caparelli J, Lee C, Pan D, Strandberg G, Vuu L and Lin Lawell C-Y C 2018 Jevons' paradox and efficient irrigation technology *Sustainability* **10** 1590
- Thomas P and Büntgen U 2017 First harvest of Périgord black truffle in the UK as a result of climate change *Clim. Res.* **74** 67–70
- Thomas P and Büntgen U 2019 A risk assessment of Europe's black truffle sector under predicted climate change *Sci. Total Environ.* **655** 27–34
- Trappe J M and Claridge A W 2010 The hidden life of truffles *Sci. Am.* **302** 78–84
- Trnka M *et al* 2018 Priority questions in multidisciplinary drought research *Clim. Res.* **75** 241–60
- Vautard R *et al* 2014 Regional climate model simulations indicate limited climatic impacts by operational and planned European wind farms *Nat. Commun.* **5** 3196
- Warren J M, Brooks J R, Meinzer F C and Eberhart J L 2008 Hydraulic redistribution of water from *pinus ponderosa* trees to seedlings: evidence for an ectomycorrhizal pathway *New Phytol.* **178** 382–94